JOURNAL OF THE AMERICAN PODIATRIC MEDICAL ASSOCIATION

January–December 2005
Volume 95

SUBJECT INDEX

A

African populations
Arch index as a predictor of pes planus: a comparative study of indigenous Kenyans and Tanzanians [Igbigbi et al], 3:273

Anatomy
A new concept of indexing tibiofibular torsion: a pilot study using dry bones [Tamari et al], 5:481

Anesthesia
New nail block technique [Ouzounov], 6:589

Ankle
Arthroscopy-assisted retrograde drilling of osteochondral lesions of the talar dome [Nelson and Haycock], 1:91
Double calcaneal osteotomy: realignment considerations in eight patients [Catanzariti et al], 1:53
Endoscopic gastrocnemius recession for the treatment of gastrocnemius equinus [DiDomenico et al], 4:410
Equinus deformity as a factor in forefoot nerve entrapment: treatment with endoscopic gastrocnemius recession [Barrett and Jarvis], 5:464
Gastrocnemius soleus recession: a simpler, more limited approach [Lamm et al], 1:18
Percutaneous supramalleolar osteotomy for distal tibial (near articular) ankle deformities [Mendicino et al], 1:72
Percutaneous tendo Achillis lengthening to promote healing of diabetic plantar foot ulceration [Willrich et al], 3:281
Realignment arthrodesis of the rearfoot and ankle: a comprehensive evaluation [Mendicino et al], 1:60
Reliability of ankle goniometric measurements: a literature review [Martin and McPoil], 6:564
Sinus tarsi denervation: clinical results [Dellon and Barrett], 2:108
A systematic approach to evaluation of the rearfoot, ankle, and leg in reconstructive surgery [Mendicino et al], 1:2

Antiseptic
A critical discussion of the use of antiseptics in acute traumatic wounds [Smith], 2:148

APMA Announcements and Reports
American Podiatric Medical Association Best Walking City Competition, 2004 [Fisher et al], 4:414

Arch index
Arch index as a predictor of pes planus: a comparative study of indigenous Kenyans and Tanzanians [Igbigbi et al], 3:273

Artery
The challenge of peripheral arterial disease in people with diabetes: a call to action [Sanders], 3:307
Peripheral arterial disease in people with diabetes [American Diabetes Association], 3:309

Arthritis
Determinants of disabling foot pain in retirement village residents [Menz and Morris], 6:573
Effect of regular exercise on the radiographic progression of foot osteoarthritis [Wilder et al], 4:342
Use of laterally wedged custom foot orthoses to reduce pain associated with medial knee osteoarthritis: a preliminary investigation [Rubin and Menz], 4:347

Arthrodesis
A new fixation technique for the Lapidus bunionectomy [Wang and Riley], 4:405; (letter) [Strash], 6:606; (author's response) [Wang], 6:607
Realignment arthrodesis of the rearfoot and ankle: a comprehensive evaluation [Mendicino et al], 1:60
Realignment considerations in the triple arthrodesis [Catanzariti et al], 1:13
Subtalar joint arthrodesis [Catanzariti et al], 1:34
Technical considerations in tarsometatarsal joint arthrodesis [Catanzariti and Mendicino], 1:85

Arthrodiastasis
Management of hallux limitus with distraction of the first metatarsophalangeal joint [Talarico et al], 2:121

Arthroscopy
Arthroscopy-assisted retrograde drilling of osteochondral lesions of the talar dome [Nelson and Haycock], 1:91

Atherosclerosis
The challenge of peripheral arterial disease in people with diabetes: a call to action [Sanders], 3:307
Peripheral arterial disease in people with diabetes [American Diabetes Association], 3:309

Authorship
An analysis of articles published in JAPMA in 2004 (letter) [Schlefman], 3:320; (editor's response) [Joseph], 3:320

B

Best Walking City Competition
American Podiatric Medical Association Best Walking City Competition, 2004 [Fisher et al], 4:414

Biomechanics
The amount of rearfoot motion used during the stance phase of walking [Youberg et al], 4:376
Effectiveness of low-Dye taping for the short-term management of plantar fasciitis [Landorf et al], 6:525
Effects of simulated genu valgum and genu varum on ground reaction forces and subtalar joint function during gait [Van Gheluwe et al], 6:531
Functional hallux limitus and lesser-metatarsal overload [Clough], 6:593
Future of gait analysis: a podiatric medical perspective [Curran and Dananberg], 2:130
Influence of treadmill design on rearfoot pronation during gait at different speeds [Sajko and Pierrynowski], 5:475

A new concept of indexing tibiofibular torsion: a pilot study using dry bones [Tamari et al], 5:481
Plantar pressure and radiographic changes to the forefoot after the Austin bunionectomy [Bryant et al], 4:357
Reliability and validity of clinically assessing first-ray mobility of the foot (letter) [Allen], 3:322; (author's response) [Cornwall], 3:322
Reliability of ankle goniometric measurements: a literature review [Martin and McPoil], 6:564
The stance phase of walking during late pregnancy: temporospatial and ground reaction force variables [Lymbery and Gilleard], 3:247
Three-dimensional reconstruction of magnetic resonance images of a displaced flexor hallucis longus tendon in hallux valgus [Sanders et al], 4:401
Use of the longitudinal arch angle to predict dynamic foot posture in walking [McPoil and Cornwall], 2:114

Blood
Low-molecular-weight heparins: an overview for the podiatric physician [Smith], 4:383
Protein C deficiency: podiatric medical relevance and case report [Motley and Vanlandingham], 5:491
Xanthoma of the calcaneus associated with hyperlipoproteinemia [Yalcinkaya and Oztop], 6:602

Bone
Arthroscopy-assisted retrograde drilling of osteochondral lesions of the talar dome [Nelson and Haycock], 1:91
Bone matrix therapy for aneurysmal bone cysts [Carpenter and Motley], 4:394
Brucella osteomyelitis of the calcaneus [Baktiroglu et al], 2:216
Callus formation of the tibial hallux sesamoid mimicking a bone tumor [Kiter et al], 6:580
Determinants of disabling foot pain in retirement village residents [Menz and Morris], 6:573
Dorsal foot pain due to compression of the deep peroneal nerve by exostosis of the metatarsocuneiform joint [Parker], 5:455
Double calcaneal osteotomy: realignment considerations in eight patients [Catanzariti et al], 1:53
Effect of regular exercise on the radiographic progression of foot osteoarthritis [Wilder et al], 4:342
Focal tuberculous osteomyelitis of the calcaneus secondary to direct extension from an infected retrocalcaneal bursa [Abdelwahab et al], 3:285
Hallucal sesamoid osteonecrosis: an overlooked cause of forefoot pain [Özkoç et al], 3:277
Melorheostosis in a pediatric patient [Schreck], 2:167
A new concept of indexing tibiofibular torsion: a pilot study using dry bones [Tamari et al], 5:481
Os vesalianum pedis [Boya et al], 6:583
Osteochondroma of the talar neck: a rare cause of callosity of the foot dorsum [Keser and Bayar], 3:295
Osteochondroma of the talus in a male adolescent [Joshi et al], 5:494
Percutaneous supramalleolar osteotomy for distal tibial (near articular) ankle deformities [Mendicino et al], 1:72
Use of laterally wedged custom foot orthoses to reduce pain associated with medial knee osteoarthritis: a preliminary investigation [Rubin and Menz], 4:347
Xanthoma of the calcaneus associated with hyperlipoproteinemia [Yalcinkaya and Oztop], 6:602

Book reviews
Atlas of Foot and Ankle Sonography (Adler et al, eds) [Stone], 6:608
An Atlas of Foot and Ankle Surgery, 2nd Ed (Wülker et al, eds) [Roukis], 6:608

Bristol Foot Score
The Bristol Foot Score: developing a patient-based foot-health measure [Barnett et al], 3:264

Brucellosis
Brucella osteomyelitis of the calcaneus [Baktiroglu et al], 2:216

Bunionectomy
Capsular interposition for the Keller bunionectomy with the use of soft-tissue anchors [Cook], 2:180
A new fixation technique for the Lapidus bunionectomy [Wang and Riley], 4:405; (letter) [Strash], 6:606; (author's response) [Wang], 6:607
Plantar pressure and radiographic changes to the forefoot after the Austin bunionectomy [Bryant et al], 4:357

Bursa
Focal tuberculous osteomyelitis of the calcaneus secondary to direct extension from an infected retrocalcaneal bursa [Abdelwahab et al], 3:285

C

Calcaneus
Brucella osteomyelitis of the calcaneus [Baktiroglu et al], 2:216
Double calcaneal osteotomy: realignment considerations in eight patients [Catanzariti et al], 1:53
Effectiveness of extracorporeal shockwave treatment in 353 patients with chronic plantar fasciitis [Norris et al], 6:517
Effectiveness of low-Dye taping for the short-term management of plantar fasciitis [Landorf et al], 6:525
Focal tuberculous osteomyelitis of the calcaneus secondary to direct extension from an infected retrocalcaneal bursa [Abdelwahab et al], 3:285
Partial calcanectomy for the treatment of recalcitrant heel ulcerations [Randall et al], 4:335
Xanthoma of the calcaneus associated with hyperlipoproteinemia [Yalcinkaya and Oztop], 6:602

Callus
Callus formation of the tibial hallux sesamoid mimicking a bone tumor [Kiter et al], 6:580
Osteochondroma of the talar neck: a rare cause of callosity of the foot dorsum [Keser and Bayar], 3:295

Cartilage
Osteochondroma of the talar neck: a rare cause of callosity of the foot dorsum [Keser and Bayar], 3:295
Osteochondroma of the talus in a male adolescent [Joshi et al], 5:494

Charcot's foot
Evaluation of Young's modulus in Achilles tendons with diabetic neuroarthropathy [Grant et al], 3:242

Cimetidine
Cimetidine as a first-line therapy for pedal verruca: eight-year retrospective analysis [Mullen et al], 3:229; (letter) [Turlik], 5:508; (authors' response) [Mullen et al], 5:508

Coagulation
Protein C deficiency: podiatric medical relevance and case report [Motley and Vanlandingham], 5:491

Cutaneous larva migrans
Cutaneous larva migrans: case report with current recommendations for treatment [O'Quinn and Dushin], 3:291

Cyst
Bone matrix therapy for aneurysmal bone cysts [Carpenter and Motley], 4:394
Tarsal tunnel surgery for treatment of tarsal ganglion: a rewarding operation with devastating potential complications [Rosson et al], 5:459

D

Deep venous thrombosis
Low-molecular-weight heparins: an overview for the podiatric physician [Smith], 4:383

Diabetes mellitus
The challenge of peripheral arterial disease in people with diabetes: a call to action [Sanders], 3:307
Cooling the foot to prevent diabetic foot wounds: a proof-of-concept trial [Armstrong et al], 2:103
Diagnosis and treatment of diabetic foot infections [Lipsky et al], 2:183
Efficacy of fifth metatarsal head resection for treatment of chronic diabetic foot ulceration [Armstrong et al], 4:353
Evaluation of Young's modulus in Achilles tendons with diabetic neuroarthropathy [Grant et al], 3:242
Improved sensitivity in patients with peripheral neuropathy: effects of monochromatic infrared photo energy [DeLellis et al], 2:143
Maggot therapy in "lower-extremity hospice" wound care: fewer amputations and more antibiotic-free days [Armstrong et al], 3:254
Morphological and functional changes in the diabetic peripheral nerve: using diagnostic ultrasound and neurosensory testing to select candidates for nerve decompression [Lee and Dauphinée], 5:433
Percutaneous tendo Achillis lengthening to promote healing of diabetic plantar foot ulceration [Willrich et al], 3:281
Peripheral arterial disease in people with diabetes [American Diabetes Association], 3:309
Surgical decompression in lower-extremity diabetic peripheral neuropathy [Rader], 5:446
Surgical treatment of peripheral neuropathy: outcomes from 100 consecutive decompressions [Valdivia et al], 5:451
Testing for loss of protective sensation in patients with foot ulceration: a cross-sectional study [Wood et al], 5:469

E

Education
Internet resources for podiatric medical students: an update [Fikar], 2:211
Podiatric medical students' evaluation of a computerized student-patient encounter log system [Morewitz et al], 6:556
A train-the-trainer model for integrating evidence-based medicine training into podiatric medical education [Green], 5:497

Equinus deformity
Endoscopic gastrocnemius recession for the treatment of gastrocnemius equinus [DiDomenico et al], 4:410
Equinus deformity as a factor in forefoot nerve entrapment: treatment with endoscopic gastrocnemius recession [Barrett and Jarvis], 5:464
Gastrocnemius soleus recession: a simpler, more limited approach [Lamm et al], 1:18
Percutaneous tendo Achillis lengthening to promote healing of diabetic plantar foot ulceration [Willrich et al], 3:281

Evidence-based medicine
A train-the-trainer model for integrating evidence-based medicine training into podiatric medical education [Green], 5:497

Exercise
Effect of regular exercise on the radiographic progression of foot osteoarthritis [Wilder et al], 4:342

Exostosis
Dorsal foot pain due to compression of the deep peroneal nerve by exostosis of the metatarsocuneiform joint [Parker], 5:455

Extracorporeal shockwave treatment
Effectiveness of extracorporeal shockwave treatment in 353 patients with chronic plantar fasciitis [Norris et al], 6:517

F

Flatfoot
Arch index as a predictor of pes planus: a comparative study of indigenous Kenyans and Tanzanians [Igbigbi et al], 3:273
Double calcaneal osteotomy: realignment considerations in eight patients [Catanzariti et al], 1:53
A simple intraoperative technique to accurately align the rearfoot complex [Roukis and Kirby], 5:505

Fluorouracil
Treatment of verruca plantaris with a combination of topical fluorouracil and salicylic acid [Young and Cohen], 4:366

Foot type
Arch index as a predictor of pes planus: a comparative study of indigenous Kenyans and Tanzanians [Igbigbi et al], 3:273
Foot type and overuse injury in triathletes [Burns et al], 3:235

Footdrop
A simple technique for correcting footdrop: suspension tenodesis of the tibialis anterior tendon to the distal tibia [Roukis et al], 2:154

Forefoot
Equinus deformity as a factor in forefoot nerve entrapment: treatment with endoscopic gastrocnemius recession [Barrett and Jarvis], 5:464
Use of high-resolution ultrasound in evaluation of the forefoot to differentiate forefoot nerve entrapments [Kincaid and Barrett], 5:429

G

Gait
The amount of rearfoot motion used during the stance phase of walking [Youberg et al], 4:376
Effects of simulated genu valgum and genu varum on ground reaction forces and subtalar joint function during gait [Van Gheluwe et al], 6:531
Functional hallux limitus and lesser-metatarsal overload [Clough], 6:593
Future of gait analysis: a podiatric medical perspective [Curran and Dananberg], 2:130
Influence of treadmill design on rearfoot pronation during gait at different speeds [Sajko and Pierrynowski], 5:475

The stance phase of walking during late pregnancy: temporospatial and ground reaction force variables [Lymbery and Gilleard], 3:247
Use of the longitudinal arch angle to predict dynamic foot posture in walking [McPoil and Cornwall], 2:114

Ganglion
Tarsal tunnel surgery for treatment of tarsal ganglion: a rewarding operation with devastating potential complications [Rosson et al], 5:459

Gangrene
Protein C deficiency: podiatric medical relevance and case report [Motley and Vanlandingham], 5:491

Gastrocnemius soleus complex
Endoscopic gastrocnemius recession for the treatment of gastrocnemius equinus [DiDomenico et al], 4:410
Equinus deformity as a factor in forefoot nerve entrapment: treatment with endoscopic gastrocnemius recession [Barrett and Jarvis], 5:464
Gastrocnemius soleus recession: a simpler, more limited approach [Lamm et al], 1:18
Percutaneous tendo Achillis lengthening to promote healing of diabetic plantar foot ulceration [Willrich et al], 3:281

Geriatric population
Determinants of disabling foot pain in retirement village residents [Menz and Morris], 6:573

Gigantism
A neglected case of macrodystrophia lipomatosa of the foot in an elderly man [Tuzuner et al], 5:486

Goniometry
Reliability of ankle goniometric measurements: a literature review [Martin and McPoil], 6:564

Guidelines, authors'
Guidelines for authors, 1:97, 3:323, 5:510

H

Hallux
Callus formation of the tibial hallux sesamoid mimicking a bone tumor [Kiter et al], 6:580
Capsular interposition for the Keller bunionectomy with the use of soft-tissue anchors [Cook], 2:180
Functional hallux limitus and lesser-metatarsal overload [Clough], 6:593
Hallucal sesamoid osteonecrosis: an overlooked cause of forefoot pain [Özkoç et al], 3:277
Management of hallux limitus with distraction of the first metatarsophalangeal joint [Talarico et al], 2:121
Metatarsus primus elevatus in hallux rigidus: fact or fiction? [Roukis], 3:221
A new fixation technique for the Lapidus bunionectomy [Wang and Riley], 4:405; (letter) [Strash], 6:606; (author's response) [Wang], 6:607
Plantar pressure and radiographic changes to the forefoot after the Austin bunionectomy [Bryant et al], 4:357
Reliability and validity of clinically assessing first-ray mobility of the foot (letter) [Allen], 3:322; (author's response) [Cornwall], 3:322
Three-dimensional reconstruction of magnetic resonance images of a displaced flexor hallucis longus tendon in hallux valgus [Sanders et al], 4:401

Hammer toe
Lesser-metatarsal medial displacement osteotomy for the treatment of digital transverse plane deformities [Goforth et al], 6:550

Heparin
Low-molecular-weight heparins: an overview for the podiatric physician [Smith], 4:383

Hookworm
Cutaneous larva migrans: case report with current recommendations for treatment [O'Quinn and Dushin], 3:291

Hyperlipoproteinemia
Xanthoma of the calcaneus associated with hyperlipoproteinemia [Yalcinkaya and Oztop], 6:602

I

Imaging
Double calcaneal osteotomy: realignment considerations in eight patients [Catanzariti et al], 1:53
Effect of regular exercise on the radiographic progression of foot osteoarthritis [Wilder et al], 4:342
Effect of tube angulation on the measurement of intermetatarsal angles [Weijers et al], 4:370
Morphological and functional changes in the diabetic peripheral nerve: using diagnostic ultrasound and neurosensory testing to select candidates for nerve decompression [Lee and Dauphinée], 5:433
Percutaneous supramalleolar osteotomy for distal tibial (near articular) ankle deformities [Mendicino et al], 1:72
Plantar pressure and radiographic changes to the forefoot after the Austin bunionectomy [Bryant et al], 4:357
Realignment arthrodesis of the rearfoot and ankle: a comprehensive evaluation [Mendicino et al], 1:60
Realignment considerations in the triple arthrodesis [Catanzariti et al], 1:13
Static rearfoot alignment: a comparison of clinical and radiographic measures [Lamm et al], 1:26
A systematic approach to evaluation of the rearfoot, ankle, and leg in reconstructive surgery [Mendicino et al], 1:2
Three-dimensional reconstruction of magnetic resonance images of a displaced flexor hallucis longus tendon in hallux valgus [Sanders et al], 4:401
Use of high-resolution ultrasound in evaluation of the forefoot to differentiate forefoot nerve entrapments [Kincaid and Barrett], 5:429

Infection
Brucella osteomyelitis of the calcaneus [Baktiroglu et al], 2:216
A comparison of KOH and culture results from two mycology laboratories for the diagnosis of onychomycosis during a randomized, multicenter clinical trial: a subset study (letter) [Arnold et al], 4:421; (authors' response) [Scherer and Scherer], 4:423
Cutaneous larva migrans: case report with current recommendations for treatment [O'Quinn and Dushin], 3:291
Diagnosis and treatment of diabetic foot infections [Lipsky et al], 2:183
Focal tuberculous osteomyelitis of the calcaneus secondary to direct extension from an infected retrocalcaneal bursa [Abdelwahab et al], 3:285
Utility of histopathologic analysis in the evaluation of onychomycosis [Weinberg et al], 3:258

Internet
Internet resources for podiatric medical students: an update [Fikar], 2:211
Podiatric medical students' evaluation of a computerized student-patient encounter log system [Morewitz et al], 6:556

J

Joint
Arthroscopy-assisted retrograde drilling of osteochondral lesions of the talar dome [Nelson and Haycock], 1:91
Capsular interposition for the Keller bunionectomy with the use of soft-tissue anchors [Cook], 2:180
Determinants of disabling foot pain in retirement village residents [Menz and Morris], 6:573
Diffuse pigmented villonodular synovitis: two cases involving the calcaneocuboid joint [Peng et al], 2:161
Dorsal foot pain due to compression of the deep peroneal nerve by exostosis of the metatarsocuneiform joint [Parker], 5:455
Double calcaneal osteotomy: realignment considerations in eight patients [Catanzariti et al], 1:53
Effect of regular exercise on the radiographic progression of foot osteoarthritis [Wilder et al], 4:342
Effects of simulated genu valgum and genu varum on ground reaction forces and subtalar joint function during gait [Van Gheluwe et al], 6:531
Endoscopic gastrocnemius recession for the treatment of gastrocnemius equinus [DiDomenico et al], 4:410
Equinus deformity as a factor in forefoot nerve entrapment: treatment with endoscopic gastrocnemius recession [Barrett and Jarvis], 5:464
Evaluation of Young's modulus in Achilles tendons with diabetic neuroarthropathy [Grant et al], 3:242
Functional hallux limitus and lesser-metatarsal overload [Clough], 6:593
Gastrocnemius soleus recession: a simpler, more limited approach [Lamm et al], 1:18
Lesser-metatarsal medial displacement osteotomy for the treatment of digital transverse plane deformities [Goforth et al], 6:550
Management of hallux limitus with distraction of the first metatarsophalangeal joint [Talarico et al], 2:121
Metatarsus primus elevatus in hallux rigidus: fact or fiction? [Roukis], 3:221
A new fixation technique for the Lapidus bunionectomy [Wang and Riley], 4:405; (letter) [Strash], 6:606; (author's response) [Wang], 6:607
Percutaneous supramalleolar osteotomy for distal tibial (near articular) ankle deformities [Mendicino et al], 1:72
Percutaneous tendo Achillis lengthening to promote healing of diabetic plantar foot ulceration [Willrich et al], 3:281
Plantar pressure and radiographic changes to the forefoot after the Austin bunionectomy [Bryant et al], 4:357
Realignment arthrodesis of the rearfoot and ankle: a comprehensive evaluation [Mendicino et al], 1:60
Realignment considerations in the triple arthrodesis [Catanzariti et al], 1:13
Reliability of ankle goniometric measurements: a literature review [Martin and McPoil], 6:564
Sinus tarsi denervation: clinical results [Dellon and Barrett], 2:108
Subtalar joint arthrodesis [Catanzariti et al], 1:34
A systematic approach to evaluation of the rearfoot, ankle, and leg in reconstructive surgery [Mendicino et al], 1:2
Technical considerations in tarsometatarsal joint arthrodesis [Catanzariti and Mendicino], 1:85
Three-dimensional reconstruction of magnetic resonance images of a displaced flexor hallucis longus tendon in hallux valgus [Sanders et al], 4:401
Use of laterally wedged custom foot orthoses to reduce pain associated with medial knee osteoarthritis: a preliminary investigation [Rubin and Menz], 4:347

K

Kaposi's sarcoma
Classic Kaposi's sarcoma [Levi], 6:586

Knee
Effects of simulated genu valgum and genu varum on ground reaction forces and subtalar joint function during gait [Van Gheluwe et al], 6:531
Use of laterally wedged custom foot orthoses to reduce pain associated with medial knee osteoarthritis: a preliminary investigation [Rubin and Menz], 4:347

L

Laser
Onychoplasty with carbon dioxide laser matrixectomy for treatment of ingrown toenails [Farley-Sakevich et al], 2:175

Lesion
Accurate nomenclature for forefoot nerve entrapment: a historical perspective [Larson et al], 3:298
Arthroscopy-assisted retrograde drilling of osteochondral lesions of the talar dome [Nelson and Haycock], 1:91
Bone matrix therapy for aneurysmal bone cysts [Carpenter and Motley], 4:394
Callus formation of the tibial hallux sesamoid mimicking a bone tumor [Kiter et al], 6:580
Cimetidine as a first-line therapy for pedal verruca: eight-year retrospective analysis [Mullen et al], 3:229; (letter) [Turlik], 5:508; (authors' response) [Mullen et al], 5:508
Classic Kaposi's sarcoma [Levi], 6:586
Diffuse pigmented villonodular synovitis: two cases involving the calcaneocuboid joint [Peng et al], 2:161
Dorsal foot pain due to compression of the deep peroneal nerve by exostosis of the metatarsocuneiform joint [Parker], 5:455
Osteochondroma of the talar neck: a rare cause of callosity of the foot dorsum [Keser and Bayar], 3:295
Osteochondroma of the talus in a male adolescent [Joshi et al], 5:494
Sinus tarsi denervation: clinical results [Dellon and Barrett], 2:108
Treatment of verruca plantaris with a combination of topical fluorouracil and salicylic acid [Young and Cohen], 4:366
Ulceration of pyoderma gangrenosum treated with negative pressure wound therapy [Geller and Longton], 2:171
Unusually large cellular schwannoma of the foot [Pasternack and Winter-Reiken], 2:157
Xanthoma of the calcaneus associated with hyperlipoproteinemia [Yalcinkaya and Oztop], 6:602

Longitudinal arch angle
Use of the longitudinal arch angle to predict dynamic foot posture in walking [McPoil and Cornwall], 2:114

M

Macrodystrophia lipomatosa
A neglected case of macrodystrophia lipomatosa of the foot in an elderly man [Tuzuner et al], 5:486

Maggot debridement therapy
Maggot therapy in "lower-extremity hospice" wound care: fewer amputations and more antibiotic-free days [Armstrong et al], 3:254

Matrixectomy
Onychoplasty with carbon dioxide laser matrixectomy for treatment of ingrown toenails [Farley-Sakevich et al], 2:175

Measurement
Effect of tube angulation on the measurement of intermetatarsal angles [Weijers et al], 4:370
Reliability of ankle goniometric measurements: a literature review [Martin and McPoil], 6:564

Melorheostosis
Melorheostosis in a pediatric patient [Schreck], 2:167

Memorial
Gerard V. Yu: 1954–2005 [Malay], 6:605

Metatarsal
Capsular interposition for the Keller bunionectomy with the use of soft-tissue anchors [Cook], 2:180
Effect of tube angulation on the measurement of intermetatarsal angles [Weijers et al], 4:370
Efficacy of fifth metatarsal head resection for treatment of chronic diabetic foot ulceration [Armstrong et al], 4:353
Functional hallux limitus and lesser-metatarsal overload [Clough], 6:593
Lesser-metatarsal medial displacement osteotomy for the treatment of digital transverse plane deformities [Goforth et al], 6:550
Metatarsus primus elevatus in hallux rigidus: fact or fiction? [Roukis], 3:221
A new fixation technique for the Lapidus bunionectomy [Wang and Riley], 4:405; (letter) [Strash], 6:606; (author's response) [Wang], 6:607
Os vesalianum pedis [Boya et al], 6:583
Plantar pressure and radiographic changes to the forefoot after the Austin bunionectomy [Bryant et al], 4:357
Three-dimensional reconstruction of magnetic resonance images of a displaced flexor hallucis longus tendon in hallux valgus [Sanders et al], 4:401

Monochromatic infrared photo energy
Improved sensitivity in patients with peripheral neuropathy: effects of monochromatic infrared photo energy [DeLellis et al], 2:143

N

Nail
A comparison of KOH and culture results from two mycology laboratories for the diagnosis of onychomycosis during a randomized, multicenter clinical trial: a subset study (letter) [Arnold et al], 4:421; (authors' response) [Scherer and Scherer], 4:423
Congenital malalignment of the great toenails in a pair of monozygotic twins [Özdemir et al], 4:398
New nail block technique [Ouzounov], 6:589
Onychoplasty with carbon dioxide laser matrixectomy for treatment of ingrown toenails [Farley-Sakevich et al], 2:175
Treatment of ingrown toenails using a new conservative method: a prospective study comparing brace treatment with Emmert's procedure [Harrer et al], 6:542
Utility of histopathologic analysis in the evaluation of onychomycosis [Weinberg et al], 3:258

Necrosis, avascular
Hallucal sesamoid osteonecrosis: an overlooked cause of forefoot pain [Özkoç et al], 3:277

Negative pressure wound therapy
Ulceration of pyoderma gangrenosum treated with negative pressure wound therapy [Geller and Longton], 2:171

Nerve
Accurate nomenclature for forefoot nerve entrapment: a historical perspective [Larson et al], 3:298
Comparison of measures of large-fiber nerve function in patients with chronic nerve compression and neuropathy [Radoiu et al], 5:438
Dorsal foot pain due to compression of the deep peroneal nerve by exostosis of the metatarsocuneiform joint [Parker], 5:455
Equinus deformity as a factor in forefoot nerve entrapment: treatment with endoscopic gastrocnemius recession [Barrett and Jarvis], 5:464
Evaluation of Young's modulus in Achilles tendons with diabetic neuroarthropathy [Grant et al], 3:242
Morphological and functional changes in the diabetic peripheral nerve: using diagnostic ultrasound and neurosensory testing to select candidates for nerve decompression [Lee and Dauphinée], 5:433
New nail block technique [Ouzounov], 6:589
Sinus tarsi denervation: clinical results [Dellon and Barrett], 2:108
Surgical decompression in lower-extremity diabetic peripheral neuropathy [Rader], 5:446
Surgical treatment of peripheral neuropathy: outcomes from 100 consecutive decompressions [Valdivia et al], 5:451
Tarsal tunnel surgery for treatment of tarsal ganglion: a rewarding operation with devastating potential complications [Rosson et al], 5:459
Testing for loss of protective sensation in patients with foot ulceration: a cross-sectional study [Wood et al], 5:469
Unusually large cellular schwannoma of the foot [Pasternack and Winter-Reiken], 2:157
Use of high-resolution ultrasound in evaluation of the forefoot to differentiate forefoot nerve entrapments [Kincaid and Barrett], 5:429

Neuroarthropathy
Evaluation of Young's modulus in Achilles tendons with diabetic neuroarthropathy [Grant et al], 3:242

Neuroma
Accurate nomenclature for forefoot nerve entrapment: a historical perspective [Larson et al], 3:298
Sinus tarsi denervation: clinical results [Dellon and Barrett], 2:108

Neuropathy
Comparison of measures of large-fiber nerve function in patients with chronic nerve compression and neuropathy [Radoiu et al], 5:438
Cooling the foot to prevent diabetic foot wounds: a proof-of-concept trial [Armstrong et al], 2:103
Diagnosis and treatment of diabetic foot infections [Lipsky et al], 2:183

Efficacy of fifth metatarsal head resection for treatment of chronic diabetic foot ulceration [Armstrong et al], 4:353
Evaluation of Young's modulus in Achilles tendons with diabetic neuroarthropathy [Grant et al], 3:242
Improved sensitivity in patients with peripheral neuropathy: effects of monochromatic infrared photo energy [DeLellis et al], 2:143
Morphological and functional changes in the diabetic peripheral nerve: using diagnostic ultrasound and neurosensory testing to select candidates for nerve decompression [Lee and Dauphinée], 5:433
Percutaneous tendo Achillis lengthening to promote healing of diabetic plantar foot ulceration [Willrich et al], 3:281
Surgical decompression in lower-extremity diabetic peripheral neuropathy [Rader], 5:446
Surgical treatment of peripheral neuropathy: outcomes from 100 consecutive decompressions [Valdivia et al], 5:451
Testing for loss of protective sensation in patients with foot ulceration: a cross-sectional study [Wood et al], 5:469

Nomenclature
Accurate nomenclature for forefoot nerve entrapment: a historical perspective [Larson et al], 3:298

O

Onychocryptosis
Congenital malalignment of the great toenails in a pair of monozygotic twins [Özdemir et al], 4:398
Onychoplasty with carbon dioxide laser matrixectomy for treatment of ingrown toenails [Farley-Sakevich et al], 2:175
Treatment of ingrown toenails using a new conservative method: a prospective study comparing brace treatment with Emmert's procedure [Harrer et al], 6:542

Onychomycosis
A comparison of KOH and culture results from two mycology laboratories for the diagnosis of onychomycosis during a randomized, multicenter clinical trial: a subset study (letter) [Arnold et al], 4:421; (authors' response) [Scherer and Scherer], 4:423
Utility of histopathologic analysis in the evaluation of onychomycosis [Weinberg et al], 3:258

Onychoplasty
Onychoplasty with carbon dioxide laser matrixectomy for treatment of ingrown toenails [Farley-Sakevich et al], 2:175

Orthoses
Use of laterally wedged custom foot orthoses to reduce pain associated with medial knee osteoarthritis: a preliminary investigation [Rubin and Menz], 4:347

Osteoarthritis
Determinants of disabling foot pain in retirement village residents [Menz and Morris], 6:573
Effect of regular exercise on the radiographic progression of foot osteoarthritis [Wilder et al], 4:342
Use of laterally wedged custom foot orthoses to reduce pain associated with medial knee osteoarthritis: a preliminary investigation [Rubin and Menz], 4:347

Osteochondroma
Osteochondroma of the talar neck: a rare cause of callosity of the foot dorsum [Keser and Bayar], 3:295
Osteochondroma of the talus in a male adolescent [Joshi et al], 5:494

Osteochondrosis
Arthroscopy-assisted retrograde drilling of osteochondral lesions of the talar dome [Nelson and Haycock], 1:91

Osteomyelitis
Brucella osteomyelitis of the calcaneus [Baktiroglu et al], 2:216
Focal tuberculous osteomyelitis of the calcaneus secondary to direct extension from an infected retrocalcaneal bursa [Abdelwahab et al], 3:285

Osteonecrosis
Hallucal sesamoid osteonecrosis: an overlooked cause of forefoot pain [Özkoç et al], 3:277

Osteotomy
Double calcaneal osteotomy: realignment considerations in eight patients [Catanzariti et al], 1:53
Lesser-metatarsal medial displacement osteotomy for the treatment of digital transverse plane deformities [Goforth et al], 6:550
Percutaneous supramalleolar osteotomy for distal tibial (near articular) ankle deformities [Mendicino et al], 1:72

Overuse injury
Foot type and overuse injury in triathletes [Burns et al], 3:235

P

Pain
Determinants of disabling foot pain in retirement village residents [Menz and Morris], 6:573

Pediatric population
Melorheostosis in a pediatric patient [Schreck], 2:167

Peripheral arterial disease
The challenge of peripheral arterial disease in people with diabetes: a call to action [Sanders], 3:307
Peripheral arterial disease in people with diabetes [American Diabetes Association], 3:309

Pes cavovarus
A simple intraoperative technique to accurately align the rearfoot complex [Roukis and Kirby], 5:505

Pes cavus
Pes cavus [Statler and Tullis], 1:42

Pes planus
Arch index as a predictor of pes planus: a comparative study of indigenous Kenyans and Tanzanians [Igbigbi et al], 3:273
Double calcaneal osteotomy: realignment considerations in eight patients [Catanzariti et al], 1:53
A simple intraoperative technique to accurately align the rearfoot complex [Roukis and Kirby], 5:505

Pharmacology
A critical discussion of the use of antiseptics in acute traumatic wounds [Smith], 2:148
Low-molecular-weight heparins: an overview for the podiatric physician [Smith], 4:383

Plantar fasciitis
Effectiveness of extracorporeal shockwave treatment in 353 patients with chronic plantar fasciitis [Norris et al], 6:517
Effectiveness of low-Dye taping for the short-term management of plantar fasciitis [Landorf et al], 6:525

Pregnancy
The stance phase of walking during late pregnancy: temporospatial and ground reaction force variables [Lymbery and Gilleard], 3:247

President's Message
Acceptance address: prospering with a new spirit of cooperation [Glickman], 3:331
Address to APMA's House of Delegates [Smith], 3:328
How Health Policy and Practice affects your practice [Smith], 2:218
Is there integrity in the CME process? [Smith], 1:102
Our amazing Annual Scientific Meeting [Glickman], 4:425
The value of corporate support [Glickman], 6:611
What 19 percent of APMA members have achieved [Glickman], 5:515

Pressure, plantar
Efficacy of fifth metatarsal head resection for treatment of chronic diabetic foot ulceration [Armstrong et al], 4:353
Partial calcanectomy for the treatment of recalcitrant heel ulcerations [Randall et al], 4:335
Percutaneous tendo Achillis lengthening to promote healing of diabetic plantar foot ulceration [Willrich et al], 3:281
Plantar pressure and radiographic changes to the forefoot after the Austin bunionectomy [Bryant et al], 4:357

Protein C
Protein C deficiency: podiatric medical relevance and case report [Motley and Vanlandingham], 5:491

Pyoderma gangrenosum
Ulceration of pyoderma gangrenosum treated with negative pressure wound therapy [Geller and Longton], 2:171

R

Rearfoot
The amount of rearfoot motion used during the stance phase of walking [Youberg et al], 4:376
Influence of treadmill design on rearfoot pronation during gait at different speeds [Sajko and Pierrynowski], 5:475
Percutaneous supramalleolar osteotomy for distal tibial (near articular) ankle deformities [Mendicino et al], 1:72
Realignment arthrodesis of the rearfoot and ankle: a comprehensive evaluation [Mendicino et al], 1:60
Realignment considerations in the triple arthrodesis [Catanzariti et al], 1:13
A simple intraoperative technique to accurately align the rearfoot complex [Roukis and Kirby], 5:505
Static rearfoot alignment: a comparison of clinical and radiographic measures [Lamm et al], 1:26
Subtalar joint arthrodesis [Catanzariti et al], 1:34
A systematic approach to evaluation of the rearfoot, ankle, and leg in reconstructive surgery [Mendicino et al], 1:2

S

Salicylic acid
Treatment of verruca plantaris with a combination of topical fluorouracil and salicylic acid [Young and Cohen], 4:366

Sarcoma
Classic Kaposi's sarcoma [Levi], 6:586

Schwannoma
Unusually large cellular schwannoma of the foot [Pasternack and Winter-Reiken], 2:157

Sesamoid
Callus formation of the tibial hallux sesamoid mimicking a bone tumor [Kiter et al], 6:580
Hallucal sesamoid osteonecrosis: an overlooked cause of forefoot pain [Özkoç et al], 3:277

Sinus tarsi
Sinus tarsi denervation: clinical results [Dellon and Barrett], 2:108

Skin
Cimetidine as a first-line therapy for pedal verruca: eight-year retrospective analysis [Mullen et al], 3:229; (letter) [Turlik], 5:508; (authors' response) [Mullen et al], 5:508
Classic Kaposi's sarcoma [Levi], 6:586
Cutaneous larva migrans: case report with current recommendations for treatment [O'Quinn and Dushin], 3:291
Treatment of verruca plantaris with a combination of topical fluorouracil and salicylic acid [Young and Cohen], 4:366
Ulceration of pyoderma gangrenosum treated with negative pressure wound therapy [Geller and Longton], 2:171

Sports medicine
Foot type and overuse injury in triathletes [Burns et al], 3:235

Surgery
Arthroscopy-assisted retrograde drilling of osteochondral lesions of the talar dome [Nelson and Haycock], 1:91
Capsular interposition for the Keller bunionectomy with the use of soft-tissue anchors [Cook], 2:180
Diffuse pigmented villonodular synovitis: two cases involving the calcaneocuboid joint [Peng et al], 2:161
Dorsal foot pain due to compression of the deep peroneal nerve by exostosis of the metatarsocuneiform joint [Parker], 5:455
Double calcaneal osteotomy: realignment considerations in eight patients [Catanzariti et al], 1:53
Efficacy of fifth metatarsal head resection for treatment of chronic diabetic foot ulceration [Armstrong et al], 4:353
Endoscopic gastrocnemius recession for the treatment of gastrocnemius equinus [DiDomenico et al], 4:410
Equinus deformity as a factor in forefoot nerve entrapment: treatment with endoscopic gastrocnemius recession [Barrett and Jarvis], 5:464
Gastrocnemius soleus recession: a simpler, more limited approach [Lamm et al], 1:18
Lesser-metatarsal medial displacement osteotomy for the treatment of digital transverse plane deformities [Goforth et al], 6:550
Management of hallux limitus with distraction of the first metatarsophalangeal joint [Talarico et al], 2:121
Morphological and functional changes in the diabetic peripheral nerve: using diagnostic ultrasound and neurosensory testing to select candidates for nerve decompression [Lee and Dauphinée], 5:433
A new fixation technique for the Lapidus bunionectomy [Wang and Riley], 4:405; (letter) [Strash], 6:606; (author's response) [Wang], 6:607
New nail block technique [Ouzounov], 6:589
Onychoplasty with carbon dioxide laser matrixectomy for treatment of ingrown toenails [Farley-Sakevich et al], 2:175
Partial calcanectomy for the treatment of recalcitrant heel ulcerations [Randall et al], 4:335
Percutaneous supramalleolar osteotomy for distal tibial (near articular) ankle deformities [Mendicino et al], 1:72
Percutaneous tendo Achillis lengthening to promote healing of diabetic plantar foot ulceration [Willrich et al], 3:281

Plantar pressure and radiographic changes to the forefoot after the Austin bunionectomy [Bryant et al], 4:357
Realignment arthrodesis of the rearfoot and ankle: a comprehensive evaluation [Mendicino et al], 1:60
Realignment considerations in the triple arthrodesis [Catanzariti et al], 1:13
A simple intraoperative technique to accurately align the rearfoot complex [Roukis and Kirby], 5:505
A simple technique for correcting footdrop: suspension tenodesis of the tibialis anterior tendon to the distal tibia [Roukis et al], 2:154
Sinus tarsi denervation: clinical results [Dellon and Barrett], 2:108
Subtalar joint arthrodesis [Catanzariti et al], 1:34
Surgical decompression in lower-extremity diabetic peripheral neuropathy [Rader], 5:446
Surgical treatment of peripheral neuropathy: outcomes from 100 consecutive decompressions [Valdivia et al], 5:451
A systematic approach to evaluation of the rearfoot, ankle, and leg in reconstructive surgery [Mendicino et al], 1:2
Tarsal tunnel surgery for treatment of tarsal ganglion: a rewarding operation with devastating potential complications [Rosson et al], 5:459
Technical considerations in tarsometatarsal joint arthrodesis [Catanzariti and Mendicino], 1:85
Treatment of ingrown toenails using a new conservative method: a prospective study comparing brace treatment with Emmert's procedure [Harrer et al], 6:542
Unusually large cellular schwannoma of the foot [Pasternack and Winter-Reiken], 2:157

Survey
The Bristol Foot Score: developing a patient-based foot-health measure [Barnett et al], 3:264
Podiatric medical students' evaluation of a computerized student-patient encounter log system [Morewitz et al], 6:556

Synovitis
Diffuse pigmented villonodular synovitis: two cases involving the calcaneocuboid joint [Peng et al], 2:161

T

Talus
Arthroscopy-assisted retrograde drilling of osteochondral lesions of the talar dome [Nelson and Haycock], 1:91
Osteochondroma of the talar neck: a rare cause of callosity of the foot dorsum [Keser and Bayar], 3:295
Osteochondroma of the talus in a male adolescent [Joshi et al], 5:494

Taping
Effectiveness of low-Dye taping for the short-term management of plantar fasciitis [Landorf et al], 6:525

Tarsal tunnel syndrome
Tarsal tunnel surgery for treatment of tarsal ganglion: a rewarding operation with devastating potential complications [Rosson et al], 5:459

Tendon
Evaluation of Young's modulus in Achilles tendons with diabetic neuroarthropathy [Grant et al], 3:242
Longitudinal tear of the tibialis anterior tendon [Hamilton and Ford], 4:390
Os vesalianum pedis [Boya et al], 6:583
Percutaneous tendo Achillis lengthening to promote healing of diabetic plantar foot ulceration [Willrich et al], 3:281
A simple technique for correcting footdrop: suspension tenodesis of the tibialis anterior tendon to the distal tibia [Roukis et al], 2:154
Three-dimensional reconstruction of magnetic resonance images of a displaced flexor hallucis longus tendon in hallux valgus [Sanders et al], 4:401

Tenodesis
A simple technique for correcting footdrop: suspension tenodesis of the tibialis anterior tendon to the distal tibia [Roukis et al], 2:154

Thromboembolic disease
Low-molecular-weight heparins: an overview for the podiatric physician [Smith], 4:383

Tibia
Longitudinal tear of the tibialis anterior tendon [Hamilton and Ford], 4:390
A new concept of indexing tibiofibular torsion: a pilot study using dry bones [Tamari et al], 5:481
Percutaneous supramalleolar osteotomy for distal tibial (near articular) ankle deformities [Mendicino et al], 1:72
A simple technique for correcting footdrop: suspension tenodesis of the tibialis anterior tendon to the distal tibia [Roukis et al], 2:154

Treadmill
Influence of treadmill design on rearfoot pronation during gait at different speeds [Sajko and Pierrynowski], 5:475

Tuberculosis
Focal tuberculous osteomyelitis of the calcaneus secondary to direct extension from an infected retrocalcaneal bursa [Abdelwahab et al], 3:285

Tumor
Accurate nomenclature for forefoot nerve entrapment: a historical perspective [Larson et al], 3:298
Bone matrix therapy for aneurysmal bone cysts [Carpenter and Motley], 4:394
Callus formation of the tibial hallux sesamoid mimicking a bone tumor [Kiter et al], 6:580
Classic Kaposi's sarcoma [Levi], 6:586
Diffuse pigmented villonodular synovitis: two cases involving the calcaneocuboid joint [Peng et al], 2:161
Dorsal foot pain due to compression of the deep peroneal nerve by exostosis of the metatarsocuneiform joint [Parker], 5:455
Osteochondroma of the talar neck: a rare cause of callosity of the foot dorsum [Keser and Bayar], 3:295
Osteochondroma of the talus in a male adolescent [Joshi et al], 5:494
Sinus tarsi denervation: clinical results [Dellon and Barrett], 2:108
Unusually large cellular schwannoma of the foot [Pasternack and Winter-Reiken], 2:157
Xanthoma of the calcaneus associated with hyperlipoproteinemia [Yalcinkaya and Oztop], 6:602

U

Ulcer
Cooling the foot to prevent diabetic foot wounds: a proof-of-concept trial [Armstrong et al], 2:103
A critical discussion of the use of antiseptics in acute traumatic wounds [Smith], 2:148
Diagnosis and treatment of diabetic foot infections [Lipsky et al], 2:183

Efficacy of fifth metatarsal head resection for treatment of chronic diabetic foot ulceration [Armstrong et al], 4:353
Improved sensitivity in patients with peripheral neuropathy: effects of monochromatic infrared photo energy [DeLellis et al], 2:143
Maggot therapy in "lower-extremity hospice" wound care: fewer amputations and more antibiotic-free days [Armstrong et al], 3:254
Partial calcanectomy for the treatment of recalcitrant heel ulcerations [Randall et al], 4:335
Percutaneous tendo Achillis lengthening to promote healing of diabetic plantar foot ulceration [Willrich et al], 3:281
Testing for loss of protective sensation in patients with foot ulceration: a cross-sectional study [Wood et al], 5:469
Ulceration of pyoderma gangrenosum treated with negative pressure wound therapy [Geller and Longton], 2:171

Ultrasound
Morphological and functional changes in the diabetic peripheral nerve: using diagnostic ultrasound and neurosensory testing to select candidates for nerve decompression [Lee and Dauphinée], 5:433
Use of high-resolution ultrasound in evaluation of the forefoot to differentiate forefoot nerve entrapments [Kincaid and Barrett], 5:429

V

Vein
Low-molecular-weight heparins: an overview for the podiatric physician [Smith], 4:383

Verruca
Cimetidine as a first-line therapy for pedal verruca: eight-year retrospective analysis [Mullen et al], 3:229; (letter) [Turlik], 5:508; (authors' response) [Mullen et al], 5:508
Treatment of verruca plantaris with a combination of topical fluorouracil and salicylic acid [Young and Cohen], 4:366

W

Walking
American Podiatric Medical Association Best Walking City Competition, 2004 [Fisher et al], 4:414

X

Xanthoma
Xanthoma of the calcaneus associated with hyperlipoproteinemia [Yalcinkaya and Oztop], 6:602

Y

Young's modulus
Evaluation of Young's modulus in Achilles tendons with diabetic neuroarthropathy [Grant et al], 3:242

AUTHOR INDEX

A

Abdelwahab, Ibrahim Fikry: Focal tuberculous osteomyelitis of the calcaneus secondary to direct extension from an infected retrocalcaneal bursa [Abdelwahab et al], 3:285

Abdul-Quader, Mohammed: Focal tuberculous osteomyelitis of the calcaneus secondary to direct extension from an infected retrocalcaneal bursa [Abdelwahab et al], 3:285

Adams, Heiko B.: Endoscopic gastrocnemius recession for the treatment of gastrocnemius equinus [DiDomenico et al], 4:410

Akpınar, Sercan: Hallucal sesamoid osteonecrosis: an overlooked cause of forefoot pain [Özkoç et al], 3:277

Akyol, Aynur: Congenital malalignment of the great toenails in a pair of monozygotic twins [Özdemir et al], 4:398

Allen, Mary K.: Reliability and validity of clinically assessing first-ray mobility of the foot (letter), 3:322

Alper, Murat: A neglected case of macrodystrophia lipomatosa of the foot in an elderly man [Tuzuner et al], 5:486

American Diabetes Association: Peripheral arterial disease in people with diabetes, 3:309

Andonian, Eugenia: Comparison of measures of large-fiber nerve function in patients with chronic nerve compression and neuropathy [Radoiu et al], 5:438

Angirasa, Arush K.: Percutaneous tendo Achillis lengthening to promote healing of diabetic plantar foot ulceration [Willrich et al], 3:281

Araç, Şükrü: Os vesalianum pedis [Boya et al], 6:583

Armstrong, David G.:

Cooling the foot to prevent diabetic foot wounds: a proof-of-concept trial [Armstrong et al], 2:103

Efficacy of fifth metatarsal head resection for treatment of chronic diabetic foot ulceration [Armstrong et al], 4:353

Maggot therapy in "lower-extremity hospice" wound care: fewer amputations and more antibiotic-free days [Armstrong et al], 3:254

Arnold, Brian: A comparison of KOH and culture results from two mycology laboratories for the diagnosis of onychomycosis during a randomized, multicenter clinical trial: a subset study (letter) [Arnold et al], 4:421

Axe, Timothy M.: Onychoplasty with carbon dioxide laser matrixectomy for treatment of ingrown toenails [Farley-Sakevich et al], 2:175

B

Baktiroglu, Lutfu: *Brucella* osteomyelitis of the calcaneus [Baktiroglu et al], 2:216

Barnett, Sue: The Bristol Foot Score: developing a patient-based foot-health measure [Barnett et al], 3:264

Barrett, John P., Jr.: Effect of regular exercise on the radiographic progression of foot osteoarthritis [Wilder et al], 4:342

Barrett, Stephen L.:

Accurate nomenclature for forefoot nerve entrapment: a historical perspective [Larson et al], 3:298

Equinus deformity as a factor in forefoot nerve entrapment: treatment with endoscopic gastrocnemius recession [Barrett and Jarvis], 5:464

Sinus tarsi denervation: clinical results [Dellon and Barrett], 2:108

Use of high-resolution ultrasound in evaluation of the forefoot to differentiate forefoot nerve entrapments [Kincaid and Barrett], 5:429

Battiston, Bruno: Accurate nomenclature for forefoot nerve entrapment: a historical perspective [Larson et al], 3:298

Bayar, Ahmet: Osteochondroma of the talar neck: a rare cause of callosity of the foot dorsum [Keser and Bayar], 3:295

Berendt, Anthony R.: Diagnosis and treatment of diabetic foot infections [Lipsky et al], 2:183

Bostancı, Seher: Congenital malalignment of the great toenails in a pair of monozygotic twins [Özdemir et al], 4:398

Boulton, Andrew J. M.: Maggot therapy in "lower-extremity hospice" wound care: fewer amputations and more antibiotic-free days [Armstrong et al], 3:254

Boya, Hakan: Os vesalianum pedis [Boya et al], 6:583

Briffa, Kathryn: A new concept of indexing tibiofibular torsion: a pilot study using dry bones [Tamari et al], 5:481

Bryant, Alan R.: Plantar pressure and radiographic changes to the forefoot after the Austin bunionectomy [Bryant et al], 4:357

Burke, Thomas J.: Improved sensitivity in patients with peripheral neuropathy: effects of monochromatic infrared photo energy [DeLellis et al], 2:143

Burns, Joshua: Foot type and overuse injury in triathletes [Burns et al], 3:235

C

Campbell, Rona: The Bristol Foot Score: developing a patient-based foot-health measure [Barnett et al], 3:264

Carnegie, Dale H.: Improved sensitivity in patients with peripheral neuropathy: effects of monochromatic infrared photo energy [DeLellis et al], 2:143

Carpenter, Brian: Bone matrix therapy for aneurysmal bone cysts [Carpenter and Motley], 4:394

Catanzariti, Alan R.:

Double calcaneal osteotomy: realignment considerations in eight patients [Catanzariti et al], 1:53

Percutaneous supramalleolar osteotomy for distal tibial (near articular) ankle deformities [Mendicino et al], 1:72

Realignment arthrodesis of the rearfoot and ankle: a comprehensive evaluation [Mendicino et al], 1:60

Realignment considerations in the triple arthrodesis [Catanzariti et al], 1:13

Static rearfoot alignment: a comparison of clinical and radiographic measures [Lamm et al], 1:26

Subtalar joint arthrodesis [Catanzariti et al], 1:34

A systematic approach to evaluation of the rearfoot, ankle, and leg in reconstructive surgery [Mendicino et al], 1:2

Technical considerations in tarsometatarsal joint arthrodesis [Catanzariti and Mendicino], 1:85

Clark, Joel R.: Podiatric medical students' evaluation of a computerized student-patient encounter log system [Morewitz et al], 6:556

Clough, James G.: Functional hallux limitus and lesser-metatarsal overload, 6:593

Cohen, Greg E.: Treatment of verruca plantaris with a combination of topical fluorouracil and salicylic acid [Young and Cohen], 4:366

Cohen, I. Kelman: Cooling the foot to prevent diabetic foot wounds: a proof-of-concept trial [Armstrong et al], 2:103

Cole, Joan H.: Plantar pressure and radiographic changes to the forefoot after the Austin bunionectomy [Bryant et al], 4:357

Cook, Keith D.: Capsular interposition for the Keller bunionectomy with the use of soft-tissue anchors, 2:180
Cornwall, Mark W.:
The amount of rearfoot motion used during the stance phase of walking [Youberg et al], 4:376
Reliability and validity of clinically assessing first-ray mobility of the foot (author's response), 3:322
Use of the longitudinal arch angle to predict dynamic foot posture in walking [McPoil and Cornwall], 2:114
Curran, Sarah A.: Future of gait analysis: a podiatric medical perspective [Curran and Dananberg], 2:130

D

Dananberg, Howard J.: Future of gait analysis: a podiatric medical perspective [Curran and Dananberg], 2:130
Dauphinée, Damien M.: Morphological and functional changes in the diabetic peripheral nerve: using diagnostic ultrasound and neurosensory testing to select candidates for nerve decompression [Lee and Dauphinée], 5:433
Deery, H. Gunner: Diagnosis and treatment of diabetic foot infections [Lipsky et al], 2:183
DeLellis, Salvatore L.: Improved sensitivity in patients with peripheral neuropathy: effects of monochromatic infrared photo energy [DeLellis et al], 2:143
Dellon, A. Lee:
Accurate nomenclature for forefoot nerve entrapment: a historical perspective [Larson et al], 3:298
Comparison of measures of large-fiber nerve function in patients with chronic nerve compression and neuropathy [Radoiu et al], 5:438
Sinus tarsi denervation: clinical results [Dellon and Barrett], 2:108
Surgical treatment of peripheral neuropathy: outcomes from 100 consecutive decompressions [Valdivia et al], 5:451
Tarsal tunnel surgery for treatment of tarsal ganglion: a rewarding operation with devastating potential complications [Rosson et al], 5:459
Testing for loss of protective sensation in patients with foot ulceration: a cross-sectional study [Wood et al], 5:469
DiDomenico, Lawrence A.: Endoscopic gastrocnemius recession for the treatment of gastrocnemius equinus [DiDomenico et al], 4:410
Dobbs, Bruce M.: Diffuse pigmented villonodular synovitis: two cases involving the calcaneocuboid joint [Peng et al], 2:161
Dombek, Michael F.: Subtalar joint arthrodesis [Catanzariti et al], 1:34
Dushin, Robert: Cutaneous larva migrans: case report with current recommendations for treatment [O'Quinn and Dushin], 3:291
Duzcan, Ender: Callus formation of the tibial hallux sesamoid mimicking a bone tumor [Kiter et al], 6:580

E

Eickmeier, Kimberly M.: Effectiveness of extracorporeal shockwave treatment in 353 patients with chronic plantar fasciitis [Norris et al], 6:517
Ekmekci, Pelin: Congenital malalignment of the great toenails in a pair of monozygotic twins [Özdemir et al], 4:398
Embil, John M.: Diagnosis and treatment of diabetic foot infections [Lipsky et al], 2:183

F

Farina, Edward J.: Effect of regular exercise on the radiographic progression of foot osteoarthritis [Wilder et al], 4:342
Farley-Sakevich, Tara: Onychoplasty with carbon dioxide laser matrixectomy for treatment of ingrown toenails [Farley-Sakevich et al], 2:175
Fikar, Charles R.: Internet resources for podiatric medical students: an update, 2:211
Fisher, Allan H., Jr.: American Podiatric Medical Association Best Walking City Competition, 2004 [Fisher et al], 4:414
Ford, Lawrence A.: Longitudinal tear of the tibialis anterior tendon [Hamilton and Ford], 4:390
Foreman, Eric J.: Evaluation of Young's modulus in Achilles tendons with diabetic neuroarthropathy [Grant et al], 3:242

G

Garchar, David: Endoscopic gastrocnemius recession for the treatment of gastrocnemius equinus [DiDomenico et al], 4:410
Gashi, Agim: Efficacy of fifth metatarsal head resection for treatment of chronic diabetic foot ulceration [Armstrong et al], 4:353
Geller, Stephen M.: Ulceration of pyoderma gangrenosum treated with negative pressure wound therapy [Geller and Longton], 2:171
Gilleard, Wendy: The stance phase of walking during late pregnancy: temporospatial and ground reaction force variables [Lymbery and Gilleard], 3:247
Glickman, Harold B.:
Acceptance address: prospering with a new spirit of cooperation, 3:331
Our amazing Annual Scientific Meeting, 4:425
The value of corporate support, 6:611
What 19 percent of APMA members have achieved, 5:515
Goforth, W. Preston: Lesser-metatarsal medial displacement osteotomy for the treatment of digital transverse plane deformities [Goforth et al], 6:550
Goldstein, Larry: Management of hallux limitus with distraction of the first metatarsophalangeal joint [Talarico et al], 2:121
Grady, John F.: Onychoplasty with carbon dioxide laser matrixectomy for treatment of ingrown toenails [Farley-Sakevich et al], 2:175
Grant, William P.: Evaluation of Young's modulus in Achilles tendons with diabetic neuroarthropathy [Grant et al], 3:242
Green, Michael L.: A train-the-trainer model for integrating evidence-based medicine training into podiatric medical education, 5:497
Guiliana, John V.: Cimetidine as a first-line therapy for pedal verruca: eight-year retrospective analysis [Mullen et al], 3:229; (authors' response), 5:508
Günal, İzge: Os vesalianum pedis [Boya et al], 6:583
Gürgey, Erbak: Congenital malalignment of the great toenails in a pair of monozygotic twins [Özdemir et al], 4:398

H

Haer, Amie: American Podiatric Medical Association Best Walking City Competition, 2004 [Fisher et al], 4:414
Hagman, Friso: Effects of simulated genu valgum and genu varum on ground reaction forces and subtalar joint function during gait [Van Gheluwe et al], 6:531
Hamilton, Graham A.: Longitudinal tear of the tibialis anterior tendon [Hamilton and Ford], 4:390
Hamilton, Scott A.: Testing for loss of protective sensation in patients with foot ulceration: a cross-sectional study [Wood et al], 5:469

Hannon, Patrick R.: The amount of rearfoot motion used during the stance phase of walking [Youberg et al], 4:376
Harrer, Jörg: Treatment of ingrown toenails using a new conservative method: a prospective study comparing brace treatment with Emmert's procedure [Harrer et al], 6:542
Harvey, Ian: The Bristol Foot Score: developing a patient-based foot-health measure [Barnett et al], 3:264
Haycock, Darryl M.: Arthroscopy-assisted retrograde drilling of osteochondral lesions of the talar dome [Nelson and Haycock], 1:91
Hermann, George: Focal tuberculous osteomyelitis of the calcaneus secondary to direct extension from an infected retrocalcaneal bursa [Abdelwahab et al], 3:285
Hersekli, Murat Ali: Hallucal sesamoid osteonecrosis: an overlooked cause of forefoot pain [Özkoç et al], 3:277
Herzenberg, John E.: Gastrocnemius soleus recession: a simpler, more limited approach [Lamm et al], 1:18
Hillstrom, Howard J.: Static rearfoot alignment: a comparison of clinical and radiographic measures [Lamm et al], 1:26
Hohenberger, Werner: Treatment of ingrown toenails using a new conservative method: a prospective study comparing brace treatment with Emmert's procedure [Harrer et al], 6:542

I

Ianiro, Goffredo: Partial calcanectomy for the treatment of recalcitrant heel ulcerations [Randall et al], 4:335
Igbigbi, Patrick S.: Arch index as a predictor of pes planus: a comparative study of indigenous Kenyans and Tanzanians [Igbigbi et al], 3:273
Isikan, Erdem: *Brucella* osteomyelitis of the calcaneus [Baktiroglu et al], 2:216

J

Jacobus, Dwayne A.: Evaluation of Young's modulus in Achilles tendons with diabetic neuroarthropathy [Grant et al], 3:242
Jacoby, Richard: Testing for loss of protective sensation in patients with foot ulceration: a cross-sectional study [Wood et al], 5:469
Jarvis, Jason: Equinus deformity as a factor in forefoot nerve entrapment: treatment with endoscopic gastrocnemius recession [Barrett and Jarvis], 5:464
Jennings, Maureen B.: Utility of histopathologic analysis in the evaluation of onychomycosis [Weinberg et al], 3:258
Jolley, David: Cooling the foot to prevent diabetic foot wounds: a proof-of-concept trial [Armstrong et al], 2:103
Joseph, Warren S.:
An analysis of articles published in JAPMA in 2004 (editor's response), 3:321
Diagnosis and treatment of diabetic foot infections [Lipsky et al], 2:183
Guidelines for authors, 1:97, 3:323, 5:510
Joshi, Deepak: Osteochondroma of the talus in a male adolescent [Joshi et al], 5:494

K

Karchmer, Adolf W.: Diagnosis and treatment of diabetic foot infections [Lipsky et al], 2:183
Kavak, Ayse: A neglected case of macrodystrophia lipomatosa of the foot in an elderly man [Tuzuner et al], 5:486
Kayaselçuk, Fazilet: Hallucal sesamoid osteonecrosis: an overlooked cause of forefoot pain [Özkoç et al], 3:277
Keenan, Anne-Maree:
Effectiveness of low-Dye taping for the short-term management of plantar fasciitis [Landorf et al], 6:525
Foot type and overuse injury in triathletes [Burns et al], 3:235
Kemerink, Gerrit J.: Effect of tube angulation on the measurement of intermetatarsal angles [Weijers et al], 4:370
Keser, Selçuk: Osteochondroma of the talar neck: a rare cause of callosity of the foot dorsum [Keser and Bayar], 3:295
Kessels, Alphons G. H.: Effect of tube angulation on the measurement of intermetatarsal angles [Weijers et al], 4:370
Kianifard, Farid: A comparison of KOH and culture results from two mycology laboratories for the diagnosis of onychomycosis during a randomized, multicenter clinical trial: a subset study (letter) [Arnold et al], 4:421
Kimbriel, Heather R.:
Cooling the foot to prevent diabetic foot wounds: a proof-of-concept trial [Armstrong et al], 2:103
Maggot therapy in "lower-extremity hospice" wound care: fewer amputations and more antibiotic-free days [Armstrong et al], 3:254
Kincaid, Brian R.: Use of high-resolution ultrasound in evaluation of the forefoot to differentiate forefoot nerve entrapments [Kincaid and Barrett], 5:429
King, Glenda L.:
Double calcaneal osteotomy: realignment considerations in eight patients [Catanzariti et al], 1:53
A systematic approach to evaluation of the rearfoot, ankle, and leg in reconstructive surgery [Mendicino et al], 1:2
Kirby, Kevin A.:
Effects of simulated genu valgum and genu varum on ground reaction forces and subtalar joint function during gait [Van Gheluwe et al], 6:531
A simple intraoperative technique to accurately align the rearfoot complex [Roukis and Kirby], 5:505
Kiter, Esat: Callus formation of the tibial hallux sesamoid mimicking a bone tumor [Kiter et al], 6:580
Klein, Michael J.: Focal tuberculous osteomyelitis of the calcaneus secondary to direct extension from an infected retrocalcaneal bursa [Abdelwahab et al], 3:285
Koestenblatt, Evelyn K.: Utility of histopathologic analysis in the evaluation of onychomycosis [Weinberg et al], 3:258
Kukla, Renee M.: Evaluation of Young's modulus in Achilles tendons with diabetic neuroarthropathy [Grant et al], 3:242
Kumar, Narendra: Osteochondroma of the talus in a male adolescent [Joshi et al], 5:494

L

Lal, Yadu: Osteochondroma of the talus in a male adolescent [Joshi et al], 5:494
Lamm, Bradley M.:
Gastrocnemius soleus recession: a simpler, more limited approach [Lamm et al], 1:18
Realignment arthrodesis of the rearfoot and ankle: a comprehensive evaluation [Mendicino et al], 1:60
Static rearfoot alignment: a comparison of clinical and radiographic measures [Lamm et al], 1:26
Subtalar joint arthrodesis [Catanzariti et al], 1:34
Landorf, Karl B.: Effectiveness of low-Dye taping for the short-term management of plantar fasciitis [Landorf et al], 6:525
Landsman, Adam S.: A simple technique for correcting foot-drop: suspension tenodesis of the tibialis anterior tendon to the distal tibia [Roukis et al], 2:154

Larson, Ethan E.: Accurate nomenclature for forefoot nerve entrapment: a historical perspective [Larson et al], 3:298
Lee, Doohi: Morphological and functional changes in the diabetic peripheral nerve: using diagnostic ultrasound and neurosensory testing to select candidates for nerve decompression [Lee and Dauphinée], 5:433
LeFrock, Jack L.: Diagnosis and treatment of diabetic foot infections [Lipsky et al], 2:183
Levi, Michael J.: Classic Kaposi's sarcoma, 6:586
Lew, Daniel P.: Diagnosis and treatment of diabetic foot infections [Lipsky et al], 2:183
Lipsky, Benjamin A.: Diagnosis and treatment of diabetic foot infections [Lipsky et al], 2:183
Longton, James A.: Ulceration of pyoderma gangrenosum treated with negative pressure wound therapy [Geller and Longton], 2:171
Lymbery, Janelle K.: The stance phase of walking during late pregnancy: temporospatial and ground reaction force variables [Lymbery and Gilleard], 3:247

M

Maben, Frank: Cooling the foot to prevent diabetic foot wounds: a proof-of-concept trial [Armstrong et al], 2:103
Mader, Jon T.: Diagnosis and treatment of diabetic foot infections [Lipsky et al], 2:183
Malay, D. Scot: Gerard V. Yu: 1954–2005, 6:605
Maloney, Christopher T., Jr.:
Accurate nomenclature for forefoot nerve entrapment: a historical perspective [Larson et al], 3:298
Surgical treatment of peripheral neuropathy: outcomes from 100 consecutive decompressions [Valdivia et al], 5:451
Martin, Billy R.: Maggot therapy in "lower-extremity hospice" wound care: fewer amputations and more antibiotic-free days [Armstrong et al], 3:254
Martin, RobRoy L.: Reliability of ankle goniometric measurements: a literature review [Martin and McPoil], 6:564
McDonald, Douglas K.: Lesser-metatarsal medial displacement osteotomy for the treatment of digital transverse plane deformities [Goforth et al], 6:550
McPoil, Thomas G.:
The amount of rearfoot motion used during the stance phase of walking [Youberg et al], 4:376
Reliability of ankle goniometric measurements: a literature review [Martin and McPoil], 6:564
Use of the longitudinal arch angle to predict dynamic foot posture in walking [McPoil and Cornwall], 2:114
Mendicino, Robert W.:
Double calcaneal osteotomy: realignment considerations in eight patients [Catanzariti et al], 1:53
Percutaneous supramalleolar osteotomy for distal tibial (near articular) ankle deformities [Mendicino et al], 1:72
Realignment arthrodesis of the rearfoot and ankle: a comprehensive evaluation [Mendicino et al], 1:60
Realignment considerations in the triple arthrodesis [Catanzariti et al], 1:13
Static rearfoot alignment: a comparison of clinical and radiographic measures [Lamm et al], 1:26
Subtalar joint arthrodesis [Catanzariti et al], 1:34
A systematic approach to evaluation of the rearfoot, ankle, and leg in reconstructive surgery [Mendicino et al], 1:2
Technical considerations in tarsometatarsal joint arthrodesis [Catanzariti and Mendicino], 1:85
Menn, Joseph J.: Testing for loss of protective sensation in patients with foot ulceration: a cross-sectional study [Wood et al], 5:469
Menz, Hylton B.:
Determinants of disabling foot pain in retirement village residents [Menz and Morris], 6:573
Use of laterally wedged custom foot orthoses to reduce pain associated with medial knee osteoarthritis: a preliminary investigation [Rubin and Menz], 4:347
Morewitz, Stephen J.: Podiatric medical students' evaluation of a computerized student-patient encounter log system [Morewitz et al], 6:556
Morris, Meg E.: Determinants of disabling foot pain in retirement village residents [Menz and Morris], 6:573
Motley, Travis A.:
Bone matrix therapy for aneurysmal bone cysts [Carpenter and Motley], 4:394
Protein C deficiency: podiatric medical relevance and case report [Motley and Vanlandingham], 5:491
Msamati, Boniface C.: Arch index as a predictor of pes planus: a comparative study of indigenous Kenyans and Tanzanians [Igbigbi et al], 3:273
Mullen, Barry R.: Cimetidine as a first-line therapy for pedal verruca: eight-year retrospective analysis [Mullen et al], 3:229; (authors' response), 5:508
Mullins, Samantha: Podiatric medical students' evaluation of a computerized student-patient encounter log system [Morewitz et al], 6:556

N

Neerings, Brian: Double calcaneal osteotomy: realignment considerations in eight patients [Catanzariti et al], 1:53
Nelson, Scott C.: Arthroscopy-assisted retrograde drilling of osteochondral lesions of the talar dome [Nelson and Haycock], 1:91
Nesheiwat, Fawaz: Cimetidine as a first-line therapy for pedal verruca: eight-year retrospective analysis [Mullen et al], 3:229; (authors' response), 5:508
Nixon, Brent P.:
Cooling the foot to prevent diabetic foot wounds: a proof-of-concept trial [Armstrong et al], 2:103
Maggot therapy in "lower-extremity hospice" wound care: fewer amputations and more antibiotic-free days [Armstrong et al], 3:254
Norden, Carl: Diagnosis and treatment of diabetic foot infections [Lipsky et al], 2:183
Norris, Donald M.: Effectiveness of extracorporeal shock-wave treatment in 353 patients with chronic plantar fasciitis [Norris et al], 6:517

O

O'Quinn, Jonathan C.: Cutaneous larva migrans: case report with current recommendations for treatment [O'Quinn and Dushin], 3:291
Odom, Richard D.: Lesser-metatarsal medial displacement osteotomy for the treatment of digital transverse plane deformities [Goforth et al], 6:550
Orsini, Roman C.: Subtalar joint arthrodesis [Catanzariti et al], 1:34
Ouzounov, Kaloian G.: New nail block technique, 6:589
Overbeek, Troy D.: Lesser-metatarsal medial displacement osteotomy for the treatment of digital transverse plane deformities [Goforth et al], 6:550
Özalay, Metin: Hallucal sesamoid osteonecrosis: an overlooked cause of forefoot pain [Özkoç et al], 3:277
Özcan, Özal: Os vesalianum pedis [Boya et al], 6:583
Özdemir, Esin: Congenital malalignment of the great toenails in a pair of monozygotic twins [Özdemir et al], 4:398

Özkoç, Gürkan: Hallucal sesamoid osteonecrosis: an overlooked cause of forefoot pain [Özkoç et al], 3:277
Oztop, Fikri: Xanthoma of the calcaneus associated with hyperlipoproteinemia [Yalcinkaya and Oztop], 6:602
Ozturk, Adil: *Brucella* osteomyelitis of the calcaneus [Baktiroglu et al], 2:216

P

Paley, Dror:
Gastrocnemius soleus recession: a simpler, more limited approach [Lamm et al], 1:18
Realignment arthrodesis of the rearfoot and ankle: a comprehensive evaluation [Mendicino et al], 1:60
Parker, Robert G.: Dorsal foot pain due to compression of the deep peroneal nerve by exostosis of the metatarsocuneiform joint, 5:455
Parlak, Ali Haydar: A neglected case of macrodystrophia lipomatosa of the foot in an elderly man [Tuzuner et al], 5:486
Pasternack, Warren A.: Unusually large cellular schwannoma of the foot [Pasternack and Winter-Reiken], 2:157
Patel, Khushroo E.: A simple technique for correcting footdrop: suspension tenodesis of the tibialis anterior tendon to the distal tibia [Roukis et al], 2:154
Peng, Hai-En: Diffuse pigmented villonodular synovitis: two cases involving the calcaneocuboid joint [Peng et al], 2:161
Perler, Adam D.: Management of hallux limitus with distraction of the first metatarsophalangeal joint [Talarico et al], 2:121
Petricca, Dina: A simple technique for correcting footdrop: suspension tenodesis of the tibialis anterior tendon to the distal tibia [Roukis et al], 2:154
Phillips, John: Partial calcanectomy for the treatment of recalcitrant heel ulcerations [Randall et al], 4:335
Pierrynowski, Michael R.: Influence of treadmill design on rearfoot pronation during gait at different speeds [Sajko and Pierrynowski], 5:475
Pourbagher, Ayşin: Hallucal sesamoid osteonecrosis: an overlooked cause of forefoot pain [Özkoç et al], 3:277

R

Rader, Andrew J.: Surgical decompression in lower-extremity diabetic peripheral neuropathy, 5:446
Radford, Joel A.: Effectiveness of low-Dye taping for the short-term management of plantar fasciitis [Landorf et al], 6:525
Radoiu, Horatio: Comparison of measures of large-fiber nerve function in patients with chronic nerve compression and neuropathy [Radoiu et al], 5:438
Raine, Sally: A new concept of indexing tibiofibular torsion: a pilot study using dry bones [Tamari et al], 5:481
Randall, David B.: Partial calcanectomy for the treatment of recalcitrant heel ulcerations [Randall et al], 4:335
Redmond, Anthony C.:
Effectiveness of low-Dye taping for the short-term management of plantar fasciitis [Landorf et al], 6:525
Foot type and overuse injury in triathletes [Burns et al], 3:235
Reeves, Christopher L.:
Percutaneous supramalleolar osteotomy for distal tibial (near articular) ankle deformities [Mendicino et al], 1:72
Realignment considerations in the triple arthrodesis [Catanzariti et al], 1:13
A systematic approach to evaluation of the rearfoot, ankle, and leg in reconstructive surgery [Mendicino et al], 1:2
Riley, Brendan M.: A new fixation technique for the Lapidus bunionectomy [Wang and Riley], 4:405
Roe, Tommy G.: Lesser-metatarsal medial displacement osteotomy for the treatment of digital transverse plane deformities [Goforth et al], 6:550
Rosales, Mark A.: Efficacy of fifth metatarsal head resection for treatment of chronic diabetic foot ulceration [Armstrong et al], 4:353
Rosson, Gedge D.:
Comparison of measures of large-fiber nerve function in patients with chronic nerve compression and neuropathy [Radoiu et al], 5:438
Tarsal tunnel surgery for treatment of tarsal ganglion: a rewarding operation with devastating potential complications [Rosson et al], 5:459
Roukis, Thomas S.:
Metatarsus primus elevatus in hallux rigidus: fact or fiction? 3:221
Review of *An Atlas of Foot and Ankle Surgery*, 2nd Ed (Wülker et al, eds), 6:608
A simple intraoperative technique to accurately align the rearfoot complex [Roukis and Kirby], 5:505
A simple technique for correcting footdrop: suspension tenodesis of the tibialis anterior tendon to the distal tibia [Roukis et al], 2:154
Rubin, Russel: Use of laterally wedged custom foot orthoses to reduce pain associated with medial knee osteoarthritis: a preliminary investigation [Rubin and Menz], 4:347

S

Sage, Ronald A.: Percutaneous tendo Achillis lengthening to promote healing of diabetic plantar foot ulceration [Willrich et al], 3:281
Sajko, Sandy S.: Influence of treadmill design on rearfoot pronation during gait at different speeds [Sajko and Pierrynowski], 5:475
Salas, Precious: Maggot therapy in "lower-extremity hospice" wound care: fewer amputations and more antibiotic-free days [Armstrong et al], 3:254
Saltrick, Karl R.: Subtalar joint arthrodesis [Catanzariti et al], 1:34
Sanders, Antal P.: Three-dimensional reconstruction of magnetic resonance images of a displaced flexor hallucis longus tendon in hallux valgus [Sanders et al], 4:401
Sanders, Lee J.: The challenge of peripheral arterial disease in people with diabetes: a call to action, 3:307
Sangalang, Melinda B.: Cooling the foot to prevent diabetic foot wounds: a proof-of-concept trial [Armstrong et al], 2:103
Scherer, Julia E.: American Podiatric Medical Association Best Walking City Competition, 2004 [Fisher et al], 4:414
Scherer, Michael D.: A comparison of KOH and culture results from two mycology laboratories for the diagnosis of onychomycosis during a randomized, multicenter clinical trial: a subset study (authors' response) [Scherer and Scherer], 4:423
Scherer, William P.: A comparison of KOH and culture results from two mycology laboratories for the diagnosis of onychomycosis during a randomized, multicenter clinical trial: a subset study (authors' response) [Scherer and Scherer], 4:423
Schlefman, Barbara S.: An analysis of articles published in JAPMA in 2004 (letter), 3:320
Schneider, Ignaz: Treatment of ingrown toenails using a new conservative method: a prospective study comparing brace treatment with Emmert's procedure [Harrer et al], 6:542

Schöffl, Volker: Treatment of ingrown toenails using a new conservative method: a prospective study comparing brace treatment with Emmert's procedure [Harrer et al], 6:542
Schon, Lew C.: Three-dimensional reconstruction of magnetic resonance images of a displaced flexor hallucis longus tendon in hallux valgus [Sanders et al], 4:401
Schreck, Michael A.: Melorheostosis in a pediatric patient, 2:167
Segler, Christopher P.: Diffuse pigmented villonodular synovitis: two cases involving the calcaneocuboid joint [Peng et al], 2:161
Senatore, John: Comparison of measures of large-fiber nerve function in patients with chronic nerve compression and neuropathy [Radoiu et al], 5:438
Shariff, Macfenton B.: Arch index as a predictor of pes planus: a comparative study of indigenous Kenyans and Tanzanians [Igbigbi et al], 3:273
Shaw, Graham P.: Podiatric medical students' evaluation of a computerized student-patient encounter log system [Morewitz et al], 6:556
Short, Brian: Maggot therapy in "lower-extremity hospice" wound care: fewer amputations and more antibiotic-free days [Armstrong et al], 3:254
Singh, A. K.: Osteochondroma of the talus in a male adolescent [Joshi et al], 5:494
Singh, Davindar: Osteochondroma of the talus in a male adolescent [Joshi et al], 5:494
Sirmatel, Ocal: *Brucella* osteomyelitis of the calcaneus [Baktiroglu et al], 2:216
Sloan, Marc: A simple technique for correcting footdrop: suspension tenodesis of the tibialis anterior tendon to the distal tibia [Roukis et al], 2:154
Smith, Lloyd S.:
Address to APMA's House of Delegates, 3:328
How Health Policy and Practice affects your practice, 2:218
Is there integrity in the CME process? 1:102
Smith, Robert G.:
A critical discussion of the use of antiseptics in acute traumatic wounds, 2:148
Low-molecular-weight heparins: an overview for the podiatric physician, 4:383
Snijders, Christiaan J.: Three-dimensional reconstruction of magnetic resonance images of a displaced flexor hallucis longus tendon in hallux valgus [Sanders et al], 4:401
Spinner, Robert J.: Tarsal tunnel surgery for treatment of tarsal ganglion: a rewarding operation with devastating potential complications [Rosson et al], 5:459
Statler, Trenton K.:
Pes cavus [Statler and Tullis], 1:42
Realignment arthrodesis of the rearfoot and ankle: a comprehensive evaluation [Mendicino et al], 1:60
Stavosky, James W.: Diffuse pigmented villonodular synovitis: two cases involving the calcaneocuboid joint [Peng et al], 2:161
Stone, Jeffrey A.: Review of *Atlas of Foot and Ankle Sonography* (Adler et al, eds), 6:608
Strash, Walter: A new fixation technique for the Lapidus bunionectomy (letter), 6:606

T

Talarico, Leonard M.: Management of hallux limitus with distraction of the first metatarsophalangeal joint [Talarico et al], 2:121
Tamari, Kotaro: A new concept of indexing tibiofibular torsion: a pilot study using dry bones [Tamari et al], 5:481
Tan, James S.: Diagnosis and treatment of diabetic foot infections [Lipsky et al], 2:183
Tandoğan, Reha N.:
Hallucal sesamoid osteonecrosis: an overlooked cause of forefoot pain [Özkoç et al], 3:277
Os vesalianum pedis [Boya et al], 6:583
Tavakkol, Amir: A comparison of KOH and culture results from two mycology laboratories for the diagnosis of onychomycosis during a randomized, multicenter clinical trial: a subset study (letter) [Arnold et al], 4:421
Tinley, Paul:
A new concept of indexing tibiofibular torsion: a pilot study using dry bones [Tamari et al], 5:481
Plantar pressure and radiographic changes to the forefoot after the Austin bunionectomy [Bryant et al], 4:357
Tullis, Brandon L.: Pes cavus [Statler and Tullis], 1:42
Turlik, Michael: Cimetidine as a first-line therapy for pedal verruca (letter), 5:508
Tuzuner, Tolga: A neglected case of macrodystrophia lipomatosa of the foot in an elderly man [Tuzuner et al], 5:486
Tzamaras, George P.: American Podiatric Medical Association Best Walking City Competition, 2004 [Fisher et al], 4:414

V

Valdivia, Juan M. V.: Surgical treatment of peripheral neuropathy: outcomes from 100 consecutive decompressions [Valdivia et al], 5:451
Van Gheluwe, Bart: Effects of simulated genu valgum and genu varum on ground reaction forces and subtalar joint function during gait [Van Gheluwe et al], 6:531
van Mameren, Henk: Effect of tube angulation on the measurement of intermetatarsal angles [Weijers et al], 4:370
Vanlandingham, Clint L.: Protein C deficiency: podiatric medical relevance and case report [Motley and Vanlandingham], 5:491
Vito, George R.: Management of hallux limitus with distraction of the first metatarsophalangeal joint [Talarico et al], 2:121

W

Walenkamp, Geert H. I. M.: Effect of tube angulation on the measurement of intermetatarsal angles [Weijers et al], 4:370
Wang, James C.: A new fixation technique for the Lapidus bunionectomy [Wang and Riley], 4:405; (author response), 6:607
Weijers, René E.:
Effect of tube angulation on the measurement of intermetatarsal angles [Weijers et al], 4:370
Three-dimensional reconstruction of magnetic resonance images of a displaced flexor hallucis longus tendon in hallux valgus [Sanders et al], 4:401
Weinand, Martin E.: Surgical treatment of peripheral neuropathy: outcomes from 100 consecutive decompressions [Valdivia et al], 5:451
Weinberg, Jeffrey M.: Utility of histopathologic analysis in the evaluation of onychomycosis [Weinberg et al], 3:258
Werber, Bruce R.: Effectiveness of extracorporeal shockwave treatment in 353 patients with chronic plantar fasciitis [Norris et al], 6:517
Werter, Scott A.: Testing for loss of protective sensation in patients with foot ulceration: a cross-sectional study [Wood et al], 5:469

Whitaker, Jeffrey M.: Realignment considerations in the triple arthrodesis [Catanzariti et al], 1:13
Wilder, Frances V.: Effect of regular exercise on the radiographic progression of foot osteoarthritis [Wilder et al], 4:342
Willrich, Amanda: Percutaneous tendo Achillis lengthening to promote healing of diabetic plantar foot ulceration [Willrich et al], 3:281
Wilson, Anne S.: Evaluation of Young's modulus in Achilles tendons with diabetic neuroarthropathy [Grant et al], 3:242
Winter-Reiken, Deborah J.: Unusually large cellular schwannoma of the foot [Pasternack and Winter-Reiken], 2:157
Wood, Michael A.: Testing for loss of protective sensation in patients with foot ulceration: a cross-sectional study [Wood et al], 5:469
Wood, William A.: Testing for loss of protective sensation in patients with foot ulceration: a cross-sectional study [Wood et al], 5:469

Y

Yagci, Baki: Callus formation of the tibial hallux sesamoid mimicking a bone tumor [Kiter et al], 6:580
Yalcinkaya, Ulviye: Xanthoma of the calcaneus associated with hyperlipoproteinemia [Yalcinkaya and Oztop], 6:602
Yazgan, Pelin: *Brucella* osteomyelitis of the calcaneus [Baktiroglu et al], 2:216
Youberg, Linda Dowdy: The amount of rearfoot motion used during the stance phase of walking [Youberg et al], 4:376
Young, Simon: Treatment of verruca plantaris with a combination of topical fluorouracil and salicylic acid [Young and Cohen], 4:366

Z

Zager, Emil: Onychoplasty with carbon dioxide laser matrixectomy for treatment of ingrown toenails [Farley-Sakevich et al], 2:175
Zeyrek, Fadile: *Brucella* osteomyelitis of the calcaneus [Baktiroglu et al], 2:216